arXiv:1407.2032v1 [cs.IT] 8 Jul 2014

A class of p-ary cyclic codes and their weight enumerators

Long Yu, Hongwei Liu*

School of Mathematics and Statistics, Central China Normal University, Wuhan, Hubei 430079, China

Abstract

Let m, k be positive integers such that $\frac{m}{\gcd(m,k)} \geq 3$, p be an odd prime and π be a primitive element of $\mathbb{F}_{p^m}$. Let $h_1(x)$ and $h_2(x)$ be the minimal polynomials of $-\pi^{-1}$ and $\pi^{-\frac{p^k+1}{2}}$ over $\mathbb{F}_p$, respectively. In the case of odd $\frac{m}{\gcd(m,k)}$, when k is even, $\gcd(m,k)$ is odd or when $\frac{k}{\gcd(m,k)}$ is odd, Zhou et al. in [25] obtained the weight distribution of a class of cyclic codes $\mathcal{C}$ over $\mathbb{F}_p$ with parity-check polynomial $h_1(x)h_2(x)$. In this paper, we further investigate this class of cyclic codes $\mathcal{C}$ over $\mathbb{F}_p$ in the rest case of odd $\frac{m}{\gcd(m,k)}$ and the case of even $\frac{m}{\gcd(m,k)}$. Moreover, we determine the weight distribution of cyclic codes $\mathcal{C}$.

Key Words cyclic code, exponential sum, quadratic form, weight distribution.

Mathematics Subject Classification 11T71 · 94B15

1 Introduction

Let p be an odd prime and q be a power of p. An $[n,k,d]$ linear code over the finite field $\mathbb{F}_q$ is a k-dimensional subspace of $\mathbb{F}_q^n$ with minimum Hamming distance d. Let A_i denote the number of codewords with Hamming weight i in a linear code $\mathcal{C}$ of length n. The weight enumerator of $\mathcal{C}$ is defined by

$$A_0 + A_1X + A_2X^2 + \cdots + A_nX^n, \quad \text{where } A_0 = 1.$$

The sequence $(A_0, A_1, \cdots, A_n)$ is called the weight distribution of the code $\mathcal{C}$.

A linear code $\mathcal{C}$ of length n is called cyclic if $(c_0, c_1, \cdots, c_{n-1}) \in \mathcal{C}$ implies $(c_{n-1}, c_0, \cdots, c_{n-2}) \in \mathcal{C}$. By identifying a codeword $(c_0, c_1, \cdots, c_{n-1}) \in \mathcal{C}$ with the polynomial

$$c_0 + c_1X + \cdots + c_{n-1}X^{n-1} \in \mathbb{F}_q[X]/(X^n - 1),$$

a cyclic code $\mathcal{C}$ of length n over $\mathbb{F}_q$ corresponds to an ideal of $\mathbb{F}_q[X]/(X^n-1)$. The monic generator $g(X)$ of this ideal is called the generator polynomial of $\mathcal{C}$, which satisfies that $g(X)|(X^n-1)$.

*Corresponding author.
Email addresses: hwliu@mail.ccnu.edu.cn (Hongwei Liu), longyu@mails.ccnu.edu.cn (Long Yu).

The polynomial $h(X) = (X^n - 1)/g(X)$ is referred to as the parity-check polynomial of $\mathcal{C}$ [13]. In general the weight distribution of cyclic codes are difficult to be determined and they are known only for a few special classes. There are some results on the weight distribution of cyclic codes whose duals have two or more zeros (see [2, 3, 5, 6, 10, 11, 14–26], and the references therein).

The following notations are fixed throughout this paper.

- Let p be an odd prime, m, k be positive integers, $d = \gcd(m, k)$, $s = \frac{m}{d} \geq 3$, $q = p^d$ and $q^* = (-1)^{\frac{q-1}{2}} q$.
- Let $\mathbb{F}_{p^m}$ be the finite field and $\mathbb{F}_{p^m}^* = \mathbb{F}_q \setminus \{0\}$, π be a primitive element of $\mathbb{F}_{p^m}$. For a given divisor l of m, the trace function from $\mathbb{F}_{p^m}$ to $\mathbb{F}_{p^l}$ is defined by $\mathrm{Tr}_l^m(x) = \sum_{i=0}^{\frac{m}{l}-1} x^{p^{li}}$.
- Let $v_2(j)$ denote the 2-adic valuation of integer j (i.e., the maximal power of 2 dividing j).
- Let SQ denote the set of square elements in $\mathbb{F}_{p^m}^*$, SQ_p denote the set of square elements in $\mathbb{F}_p^*$, u_p be a primitive element in $\mathbb{F}_p$, and ζ_p be a primitive p-th unity root.

Let $h_1(x)$ and $h_2(x)$ be the minimal polynomials of $-\pi^{-1}$ and $\pi^{-\frac{p^k+1}{2}}$ over $\mathbb{F}_p$, respectively. It is easy to check that $h_1(x)$ and $h_2(x)$ are polynomials of degree m and are pairwise distinct. In this paper, we let $\mathcal{C}$ be a cyclic code with parity-check polynomial $h_1(x)h_2(x)$, and then $\dim_{\mathbb{F}_p} \mathcal{C} = 2m$. By the well-known Delsarte's Theorem [1], cyclic code $\mathcal{C}$ can be expressed as

$$\mathcal{C} = \left\{ \mathbf{c}(\alpha, \beta) = \left(\mathrm{Tr}_1^m(\alpha \pi^{\frac{p^k+1}{2} i} + \beta(-\pi)^i) \right)_{i=0}^{p^m-2} \mid \alpha, \beta \in \mathbb{F}_{p^m} \right\}. \tag{1.1}$$

Recently, Zhou et al. in [25] studied this class of cyclic codes $\mathcal{C}$ in the case of odd s, when k is even, d is odd or when $\frac{k}{d}$ is odd. They showed cyclic codes $\mathcal{C}$ have only three nonzero weights. In this paper, we further investigate this class of cyclic codes $\mathcal{C}$ for the rest cases, and determine the weight distribution of $\mathcal{C}$.

The aim of this paper is to determine the weight distribution of a class of cyclic codes $\mathcal{C}$ over $\mathbb{F}_p$ defined by (1.1). To this end, by using the value distribution of the exponential sum $\sum_{x \in \mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1} + \beta x^2)}$, $\alpha, \beta \in \mathbb{F}_{p^m}$ (see Theorem 1[11]) and investigating some overdetermine equations over finite fields, we obtain the value distribution of the exponential sum $\sum_{x \in \mathbb{F}_{p^m}} (\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1} + \beta x^2)} + \zeta_p^{\mathrm{Tr}_1^m(\alpha \pi^{\frac{p^k+1}{2}} x^{p^k+1} - \beta \pi x^2)})$, $\alpha, \beta \in \mathbb{F}_{p^m}$. Applying these results, the weight distribution of cyclic codes $\mathcal{C}$ over $\mathbb{F}_p$ defined by (1.1) is obtained.

This paper is organized as follows. Section **2** gives some preliminary results. In Section **3**, we give the weight distribution of a class of cyclic codes $\mathcal{C}$ defined by (1.1) over $\mathbb{F}_p$.

2 Preliminaries

In the following, we give a brief introduction to the theory of quadratic forms over finite fields, which is needed to calculate the weight distribution of cyclic codes $\mathcal{C}$ in the next section.

Definition 2.1. *Let $\{\omega_1, \omega_2, \cdots, \omega_s\}$ be a basis for $\mathbb{F}_{q^s}$ over $\mathbb{F}_q$ and $x = \sum_{i=1}^{s} x_i\omega_i$, where $x_i \in \mathbb{F}_q$. A function $f(x)$ from $\mathbb{F}_{q^s}$ to $\mathbb{F}_q$ is called a quadratic form if it can be represented as*

$$f(x) = f(\sum_{i=1}^{s} x_i\omega_i) = \sum_{1\le i\le j\le s} a_{ij}x_ix_j, \qquad a_{ij} \in \mathbb{F}_q.$$

The rank of the quadratic form $f(x)$ is defined as the codimension of the $\mathbb{F}_q$-vector space

$$V = \{x \in \mathbb{F}_{q^s} \mid f(x+z) - f(x) - f(z) = 0, \ \text{ for all } z \in \mathbb{F}_{q^s}\},$$

denotes by rank r. Then $|V| = q^{s-r}$.

For a quadratic form $f(x)$ with s variables over $\mathbb{F}_q$, there exists a symmetric matrix A of order s over $\mathbb{F}_q$ such that $f(x) = XAX'$, where $X = (x_1, x_2, \cdots, x_s) \in \mathbb{F}_q^s$ and X' denotes the transpose of X. It is known that there exists a nonsingular matrix T over $\mathbb{F}_q$ such that TAT' is a diagonal matrix [8]. Making a nonsingular linear substitution $X = ZT$ with $Z = (z_1, z_2, \cdots, z_s) \in \mathbb{F}_q^s$, we have

$$f(x) = Z(TAT')Z' = \sum_{i=1}^{r} a_iz_i^2, \qquad a_i \in \mathbb{F}_q^*, \tag{2.1}$$

where r is the rank of $f(x)$. We have the following result (see Lemma 1 [11]).

Lemma 2.2. [11] *Let η_d be the quadratic (multiplicative) character of $\mathbb{F}_{p^d}$. For the quadratic form $f(x)$ defined by (2.1), we have*

$$\sum_{x\in\mathbb{F}_q^s} \zeta_p^{\mathrm{Tr}_1^d(f(x))} = \begin{cases} \eta_d(a_1\cdots a_r)q^{s-\frac{r}{2}}, & \text{if } q \equiv 1 \pmod 4; \\ (\sqrt{-1})^r\eta_d(a_1\cdots a_r)q^{s-\frac{r}{2}}, & \text{if } q \equiv 3 \pmod 4. \end{cases}$$

By Definition 2.1, we have that $f_{\alpha,\beta}(x) = \mathrm{Tr}_d^m(\alpha x^{p^k+1} + \beta x^2)$ is a quadratic form over $\mathbb{F}_q$. Let $r_{\alpha,\beta}$ denote the rank of $f_{\alpha,\beta}(x)$. Therefore, $r_{\alpha,\beta}$ is determined by the codimension of

$$V = \{x \in \mathbb{F}_{q^s} \mid f_{\alpha,\beta}(x+z) - f_{\alpha,\beta}(x) - f_{\alpha,\beta}(z) = 0, \ \text{ for all } z \in \mathbb{F}_{q^s}\},$$

which is determined by the number of solutions of

$$\phi_{\alpha,\beta}(x) = 0, \quad \text{where } \ \phi_{\alpha,\beta}(x) = \alpha^{p^k}x^{p^{2k}} + 2\beta^{p^k}x^{p^k} + \alpha x. \tag{2.2}$$

Hence, we have the following result (see Lemma 2 [11]).

Lemma 2.3. [11] *For $(\alpha,\beta) \in \mathbb{F}_{p^m}^2 \setminus \{(0,0)\}$, $r_{\alpha,\beta}$ is s, $s-1$ or $s-2$. Furthermore, let n_i be the number of $f_{\alpha,\beta}(x)$ with $r_{\alpha,\beta} = s-i$ for $(\alpha,\beta) \in \mathbb{F}_{p^m}^2 \setminus \{(0,0)\}$ and $i = 0,1,2$, then*

$$n_1 = \frac{(p^m-1)(p^{m-1}-1)}{p^{2d}-1}, \ n_2 = p^{m-d}(p^m-1), \ n_0 = p^{2m} - 1 - n_1 - n_2.$$

We define

$$T(\alpha,\beta) = \sum_{x\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1}+\beta x^2)}, \ \alpha,\beta \in \mathbb{F}_{p^m}. \tag{2.3}$$

Lemma 2.4. [11] *With the notations given above. The value distribution of the exponential sum $T(\alpha,\beta)$ defined by (2.3) is shown in the following:*

i) for the case s is odd,

value	*frequency*
$\pm\sqrt{q^*}q^{\frac{s-1}{2}}$,	$\frac{1}{2}p^{2d}(p^m-p^{m-d}-p^{m-2d}+1)(p^m-1)/(p^{2d}-1)$
$p^{\frac{m+d}{2}}$	$\frac{1}{2}p^{\frac{m-d}{2}}(p^{\frac{m-d}{2}}+1)(p^m-1)$
$-p^{\frac{m+d}{2}}$	$\frac{1}{2}p^{\frac{m-d}{2}}(p^{\frac{m-d}{2}}-1)(p^m-1)$
$\pm\sqrt{q^*}q^{\frac{s+1}{2}}$	$\frac{1}{2}(p^{m-d}-1)(p^m-1)/(p^{2d}-1)$
p^m	1

ii) for the case s is even,

value	*frequency*
$p^{\frac{m}{2}}$	$\frac{1}{2}p^{2d}(p^m-p^{m-d}-p^{m-2d}+p^{\frac{m}{2}}-p^{\frac{m}{2}-d}+1)(p^m-1)/(p^{2d}-1)$
$-p^{\frac{m}{2}}$	$\frac{1}{2}p^{2d}(p^m-p^{m-d}-p^{m-2d}-p^{\frac{m}{2}}+p^{\frac{m}{2}-d}+1)(p^m-1)/(p^{2d}-1)$
$\pm\sqrt{q^*}q^{\frac{s}{2}}$	$\frac{1}{2}p^{m-d}(p^m-1)$
$p^{\frac{m}{2}+d}$	$\frac{1}{2}(p^{\frac{m}{2}}-1)(p^{\frac{m}{2}-d}+1)(p^m-1)/(p^{2d}-1)$
$-p^{\frac{m}{2}+d}$	$\frac{1}{2}(p^{\frac{m}{2}}+1)(p^{\frac{m}{2}-d}-1)(p^m-1)/(p^{2d}-1)$
p^m	1

For convenience, we define $\varepsilon=\pm1$, $i=0,1,2$, $N_i=\{(\alpha,\beta)\in\mathbb{F}_q^2 \mid r_{\alpha,\beta}=s-i\}$,

$$N_{\varepsilon,i}=\begin{cases}\{(\alpha,\beta)\in\mathbb{F}_{p^m}^2\setminus\{(0,0)\} \mid T(\alpha,\beta)=\varepsilon q_0^{\frac{s+i}{2}}\}, & \text{if } s+i \text{ is even;}\\ \{(\alpha,\beta)\in\mathbb{F}_{p^m}^2\setminus\{(0,0)\} \mid T(\alpha,\beta)=\varepsilon\sqrt{q_0^*}q_0^{\frac{s+i-1}{2}}\}, & \text{if } s+i \text{ is odd,}\end{cases}$$

and $n_{\varepsilon,i}=\mid N_{\varepsilon,i}\mid$. Then, $N_i=N_{1,i}\cup N_{-1,i}$ and $n_i=n_{1,i}+n_{-1,i}$. By (2.2), we define

$$\psi(\alpha,x)=-\frac{1}{2}x^{-1}(\alpha x^{p^k}+\alpha^{p^{m-k}}x^{p^{m-k}}). \tag{2.4}$$

Hence, $\beta=\psi(\alpha,x)$ if and only if $\phi_{\alpha,\beta}(x)=0$ with $x\in\mathbb{F}_q^*$. Therefore, it is easy to see that $r_{\alpha,\beta}=s-1$ or $s-2$ if and only if, $\alpha\in\mathbb{F}_{p^m}^*$ and there exists $x\in\mathbb{F}_{p^m}^*$ such that $\beta=\psi(\alpha,x)$. Furthermore, from Lemma 2.3, we obtain the number of solutions of $\beta=\psi(\alpha,x)$ with $x\in\mathbb{F}_{p^m}^*$ is

$$\#\{x\in\mathbb{F}_{p^m}^* \mid \beta=\psi(\alpha,x)\}=\begin{cases}0, & \text{iff } (\alpha,\beta)\in N_0;\\ p^d-1, & \text{iff } (\alpha,\beta)\in N_1;\\ p^{2d}-1, & \text{iff } (\alpha,\beta)\in N_2.\end{cases} \tag{2.5}$$

At the end of this section, we investigate two classes of overdeterminate equations over finite fields.

Lemma 2.5. *Let E_1 denote the number of solutions $(x,y)\in\mathbb{F}_{p^m}^2$ of the following system equations*

$$\begin{cases}x^2+y^2=0,\\ x^{p^k+1}+y^{p^k+1}=0.\end{cases} \tag{2.6}$$

i) If $1\le v_2(m)<v_2(k)$, then $E_1=2p^m-1$.
ii) If $v_2(k)<v_2(m)$, then $E_1=2p^m-1$ for $p^k\equiv1 \pmod 4$, $E_1=1$ for $p^k\equiv3 \pmod 4$.

Proof. By the first equation of (2.6), we have $-x^2 = y^2$. Then

$$x^{p^k+1} + y^{p^k+1} = x^{p^k+1} + (-x^2)^{\frac{p^k+1}{2}} = \begin{cases} 0, & \text{if } p^k \equiv 1 \pmod 4; \\ 2x^{p^k+1}, & \text{if } p^k \equiv 3 \pmod 4. \end{cases} \tag{2.7}$$

If $1 \le v_2(m) < v_2(k)$, then m, k are both even, which implies that $p^m \equiv 1 \pmod 4$ and $p^k \equiv 1 \pmod 4$. From (2.7) and the system equations (2.6), we have E_1 is equal to the number of solutions of equation $-x^2 = y^2$, i.e., $x = \pm\pi^{\frac{p^m-1}{4}} y$. Then $E_1 = 2p^m - 1$.

If $v_2(k) < v_2(m)$, we discuss it in two cases.

Case I, when $p^k \equiv 1 \pmod 4$: E_1 is equal to the number of solutions of equation $-x^2 = y^2$ by (2.6) and (2.7). Note that m is even, then $4 \mid p^m - 1$. So, $E_1 = 2p^m - 1$.

Case II, when $p^k \equiv 3 \pmod 4$: From (2.7), we have $x^{p^k+1} + y^{p^k+1} = 2x^{p^k+1} = 0$, then $x = 0$ and $y = 0$. Hence, $E_1 = 1$. □

Lemma 2.6. *Let E_2 denote the number of solutions $(x, y, z) \in \mathbb{F}_{p^m}^3$ of the following system equations*

$$\begin{cases} x^2 + y^2 - \pi z^2 = 0, \\ x^{p^k+1} + y^{p^k+1} + \pi^{\frac{p^k+1}{2}} z^{p^k+1} = 0. \end{cases} \tag{2.8}$$

If $v_2(k) < v_2(m)$, then $E_2 = 2p^m - 1$ for $p^k \equiv 1 \pmod 4$, $E_2 = 1$ for $p^k \equiv 3 \pmod 4$.

Proof. We distinguish between the following two cases to calculate the number of solutions $(x, y, z) \in \mathbb{F}_{p^m}^3$ of (2.8).

Case I, when $z = 0$: If $z = 0$, by Lemma 2.5 ii), we get that (2.8) has $2p^m - 1$ solutions for $p^k \equiv 1 \pmod 4$ or one solution for $p^k \equiv 3 \pmod 4$.

Case II, when $z \neq 0$: It is easy to check that $x \neq 0$ and $y \neq 0$. In this case, one has that (2.8) has $(p^m - 1)M$ solutions, where M is the number of solutions of the following system equations

$$\begin{cases} x^2 + y^2 = \pi, \\ x^{p^k+1} + y^{p^k+1} = -\pi^{\frac{p^k+1}{2}}. \end{cases} \tag{2.9}$$

By (2.9), we have $(x^2 + y^2)^{p^k+1} = \pi^{p^k+1}$ and $(x^{p^k+1} + y^{p^k+1})^2 = \pi^{p^k+1}$. Combining these two equations (the first one minus the second one) leads to

$$(xy^{p^k} - x^{p^k} y)^2 = 0,$$

which shows that $x = \lambda y$, where $\lambda \in \mathbb{F}_{p^d}^*$ (since $\gcd(p^k - 1, p^m - 1) = p^d - 1$). Substituting $x = \lambda y$ into (2.9), we get

$$\begin{cases} (\lambda^2 + 1)y^2 = \pi, \\ (\lambda^2 + 1)y^{p^k+1} = -\pi^{\frac{p^k+1}{2}}. \end{cases} \tag{2.10}$$

Eliminating $\lambda^2 + 1$ of (2.10), one has

$$y^{p^k-1} = -\pi^{\frac{p^k-1}{2}} = \pi^{\frac{p^k-1}{2} + \frac{p^m-1}{2}},$$

which induces that $p^d - 1 \mid \frac{p^k-1}{2} + \frac{p^m-1}{2}$. Since $v_2(k) < v_2(m)$, then $s = m/d$ is even. Hence, $p^d - 1 \mid \frac{p^m-1}{2}$. From $v_2(k) < v_2(m)$, we have k/d is odd and then $p^d - 1 \nmid \frac{p^k-1}{2}$. It is a contradiction with $p^d - 1 \mid \frac{p^k-1}{2} + \frac{p^m-1}{2}$. So the number of solutions of (2.10) is 0, i.e., $M = 0$.

Therefore, $E_2 = 2p^m - 1$ if $p^k \equiv 1 \pmod 4$, $E_2 = 1$ if $p^k \equiv 3 \pmod 4$. □

3 The weight distribution of $\mathcal{C}$

In this section, we first calculate the weight of the codeword $\mathbf{c}(\alpha,\beta)\in\mathcal{C}$ defined by (1.1),

$$\begin{aligned}
wt(\mathbf{c}(\alpha,\beta)) &= \#\{0\le i\le p^m-2 : c_i\neq 0\}\\
&= p^m-1-\frac{1}{p}\sum_{i=0}^{p^m-2}\sum_{u\in\mathbb{F}_p}\zeta_p^{u\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}i}+\beta(-\pi)^i)}\\
&= p^m-1-\frac{1}{p}\sum_{u\in\mathbb{F}_p}\sum_{i=0}^{\frac{p^m-3}{2}}\left(\zeta_p^{u\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}2i}+\beta\pi^{2i})}+\zeta_p^{u\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}(2i+1)}-\beta\pi^{(2i+1)})}\right)\\
&= p^m-1-\frac{1}{p}\sum_{u\in\mathbb{F}_p}\sum_{x\in SQ}\left(\zeta_p^{u\mathrm{Tr}_1^m(\alpha x^{\frac{p^k+1}{2}}+\beta x)}+\zeta_p^{u\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}x^{\frac{p^k+1}{2}}-\beta\pi x)}\right)\\
&= p^m-1-\frac{1}{2p}\sum_{u\in\mathbb{F}_p}\sum_{x\in\mathbb{F}_{p^m}^*}\left(\zeta_p^{u\mathrm{Tr}_1^m(\alpha x^{p^k+1}+\beta x^2)}+\zeta_p^{u\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-\beta\pi x^2)}\right)\\
&= p^m-p^{m-1}-\frac{1}{2p}\sum_{u\in\mathbb{F}_p^*}S(u\alpha,u\beta), \qquad (3.1)
\end{aligned}$$

where

$$\begin{aligned}
S(\alpha,\beta) &= \sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1}+\beta x^2)}+\zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-\beta\pi x^2)}\right)\\
&= T(\alpha,\beta)+T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta), \qquad (3.2)
\end{aligned}$$

where $T(\alpha,\beta)$ is defined by (2.3).

From (3.1), the weight distribution of $\mathcal{C}$ is completely determined by the value distribution of $S(\alpha,\beta)$. To obtain the value distribution of $S(\alpha,\beta)$, we need a series of lemmas. Before introducing them, we define $f(x)=\mathrm{Tr}_d^m(\alpha x^{p^k+1}+\beta x^2)$ and $g(x)=\mathrm{Tr}_d^m(\pi^{\frac{p^k+1}{2}}\alpha x^{p^k+1}-\pi\beta x^2)$. Let r_f and r_g be the rank of $f(x)$ and $g(x)$, respectively. Then, we have the following result.

Lemma 3.1. *For $(\alpha,\beta)\in\mathbb{F}_{p^m}^2\setminus\{(0,0)\}$, when $1\le v_2(m)<v_2(k)$ or $v_2(k)<v_2(m)$, we have that at least one of r_f and f_g is s.*

Proof. Assume that r_f and f_g are both less than s. By (2.2), we have that there exists $x_1,x_2\in\mathbb{F}_{p^m}^*$ such that

$$\phi_{\alpha,\beta}(x_1)=\alpha^{p^k}x_1^{p^{2k}}+2\beta^{p^k}x_1^{p^k}+\alpha x_1=0, \qquad (3.3)$$

and

$$\phi_{\pi^{(p^k+1)/2}\alpha,-\pi\beta}(x_2)=(\pi^{\frac{p^k+1}{2}}\alpha)^{p^k}x_2^{p^{2k}}-2(\pi\beta)^{p^k}x_2^{p^k}+\pi^{\frac{p^k+1}{2}}\alpha x_2=0. \qquad (3.4)$$

If $\alpha=0$, we have $x_1=x_2=0$, which is a contradiction with $x_1,x_2\in\mathbb{F}_{p^m}^*$. In the following, we assume that $\alpha\neq 0$. Simplifying $(3.3)\times(\pi x_2)^{p^k}+(3.4)\times x_1^{p^k}$, one has

$$-\pi^{\frac{p^k-1}{2}}\left(\alpha x_1x_2(x_1^{p^k-1}+\pi^{\frac{p^k-1}{2}}x_2^{p^k-1})\right)^{p^k}=\alpha x_1x_2(x_1^{p^k-1}+\pi^{\frac{p^k-1}{2}}x_2^{p^k-1}), \qquad (3.5)$$

which implies that

$$x_1^{p^k-1} + \pi^{\frac{p^k-1}{2}} x_2^{p^k-1} = 0$$

or

$$\left(\alpha x_1 x_2 (x_1^{p^k-1} + \pi^{\frac{p^k-1}{2}} x_2^{p^k-1})\right)^{p^k-1} = -\pi^{-\frac{p^k-1}{2}}.$$

If one of the above two equations holds, we have that $p^d - 1 \mid \frac{p^m-1}{2} - \frac{p^k-1}{2}$.

Case I, when $1 \le v_2(m) < v_2(k)$: In this case, $s = m/d$ is odd and we get that $p^d - 1 \nmid \frac{p^m-1}{2}$. From $1 \le v_2(m) < v_2(k)$, we have that k/d is even. This implies that $p^d - 1 \mid \frac{p^k-1}{2}$ and then it is a contradiction with $p^d - 1 \mid \frac{p^m-1}{2} - \frac{p^k-1}{2}$.

Case II, $v_2(k) < v_2(m)$: Similarly, we can prove that $p^d - 1 \nmid \frac{p^m-1}{2} - \frac{p^k-1}{2}$, which is also a contradiction with $p^d - 1 \mid \frac{p^m-1}{2} - \frac{p^k-1}{2}$.

Therefore, we have that at least one of r_f and f_g is s. □

3.1 The weight distribution of $\mathcal{C}$ for $1 \le v_2(m) < v_2(k)$

In this subsection, we always assume that $1 \le v_2(m) < v_2(k)$. To determine the value distribution of $S(\alpha, \beta)$, we need the following lemma.

Lemma 3.2. *With the notations given above, we have the following result:*

1. $\sum_{\alpha,\beta \in \mathbb{F}_{p^m}} S(\alpha,\beta)^2 = 4p^{3m}$.
2. $(p^d-1) \sum_{(\alpha,\beta)\in N_1} S(\alpha,\beta)^2 + (p^{2d}-1) \sum_{(\alpha,\beta)\in N_2} S(\alpha,\beta)^2 = p^m(p^m-1)(2p^{m+d} - 2p^m + 2p^d - p^{2d} - 1)$.

Proof. 1. By (3.2), we get that

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta)^2 = \sum_{\alpha,\beta\in\mathbb{F}_{p^m}} \left(T(\alpha,\beta)^2 + 2T(\alpha,\beta)T(\pi^{\frac{p^k+1}{2}}\alpha, -\pi\beta) + T(\pi^{\frac{p^k+1}{2}}\alpha, -\pi\beta)^2 \right). \quad (3.6)$$

On one hand, by (2.3), we have

$$\begin{aligned} \sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)^2 &= \sum_{x,y\in\mathbb{F}_{p^m}} \sum_{\alpha\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha(x^{p^k+1}+y^{p^k+1}))} \sum_{\beta\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha(x^2+y^2))} \\ &= Mp^{2m}, \end{aligned} \quad (3.7)$$

where $M = \#\{(x,y) \in \mathbb{F}_{p^m}^2 \mid x^2 + y^2 = 0, x^{p^k+1} + y^{p^k+1} = 0\}$. By Lemma 2.5 i) and (3.7),

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)^2 = p^{2m}(2p^m - 1).$$

Similarly,

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\pi^{\frac{p^k+1}{2}}\alpha, -\pi\beta)^2 = p^{2m}(2p^m - 1).$$

On the other hand, by (2.3),

$$\begin{aligned}
&\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)\\
&= \sum_{x,y\in\mathbb{F}_{p^m}}\sum_{\alpha\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha(x^{p^k+1}+\pi^{\frac{p^k+1}{2}}y^{p^k+1}))}\sum_{\beta\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\beta(x^2-\pi y^2))}\\
&= \#\{(x,y)\in\mathbb{F}_{p^m}^2 \mid x^2-\pi y^2=0, x^{p^k+1}+\pi^{\frac{p^k+1}{2}}y^{p^k+1}=0\}\cdot p^{2m}\\
&= \#\{(x,y)\in\mathbb{F}_{p^m}^2 \mid x^2-\pi y^2=0, 2\pi^{\frac{p^k+1}{2}}y^{p^k+1}=0\}\cdot p^{2m}\\
&= p^{2m}.
\end{aligned}$$

Hence, from (3.6),

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta)^2 = 4p^{3m}.$$

2. By (2.5) and (3.2), we have that

$$\begin{aligned}
&(p^d-1)\sum_{(\alpha,\beta)\in N_1} S(\alpha,\beta)^2 + (p^{2d}-1)\sum_{(\alpha,\beta)\in N_2} S(\alpha,\beta)^2\\
&= \sum_{x,\alpha\in\mathbb{F}_{p^m}^*} S(\alpha,\psi(\alpha,x))^2 = \sum_{x,\alpha\in\mathbb{F}_{p^m}^*}\Big(T(\alpha,\psi(\alpha,x))+T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\psi(\alpha,x))\Big)^2. \qquad (3.8)
\end{aligned}$$

By (2.3) and (2.4),

$$\begin{aligned}
&\sum_{x,\alpha\in\mathbb{F}_{p^m}^*} T(\alpha,\psi(\alpha,x))^2\\
&= \sum_{x,\alpha\in\mathbb{F}_{p^m}^*}\sum_{y,z\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha(y^{p^k+1}+y^{p^k+1})-\frac{1}{2}(\alpha x^{p^k-1}+\alpha^{p^{m-k}}x^{p^{m-k}-1})(y^2+z^2))}\\
&= \sum_{x,\alpha\in\mathbb{F}_{p^m}^*}\sum_{y,z\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha(y^{p^k+1}+z^{p^k+1})-\frac{1}{2}\alpha x^{p^k-1}(y^2+z^2)-\frac{1}{2}x^{1-p^k}\alpha(y^{2p^k}+z^{2p^k}))}\\
&= \sum_{x\in\mathbb{F}_{p^m}^*,y,z\in\mathbb{F}_{p^m}}\sum_{\alpha\in\mathbb{F}_{p^m}^*} \zeta_p^{\mathrm{Tr}_1^m(-\frac{1}{2}\alpha x^{1-p^k}((x^{p^k-1}y-y^{p^k})^2+(x^{p^k-1}z-z^{p^k})^2))}\\
&= (p^m-1)M_1 - (p^{2m}(p^m-1)-M_1)\\
&= p^mM_1 - p^{3m}+p^{2m}, \qquad (3.9)
\end{aligned}$$

where

$$M_1 = \#\{x\in\mathbb{F}_{p^m}^*, y,z\in\mathbb{F}_{p^m} \mid (x^{p^k-1}y-y^{p^k})^2+(x^{p^k-1}z-z^{p^k})^2=0\}.$$

For a fixed $x\in\mathbb{F}_{p^m}^*$, we investigate the following equation

$$(x^{p^k-1}y-y^{p^k})^2 = -(x^{p^k-1}z-z^{p^k})^2. \qquad (3.10)$$

Since m is even, then $4 \mid p^m-1$. This shows that $-1=\pi^{\frac{p^m-1}{2}}$ is a square element in $\mathbb{F}_{p^m}$.

In the following, we discuss it case by case.

Case I, when $y=0$: In this case, we have $x^{p^k-1}z-z^{p^k}=0$, i.e., $z=\lambda x$, where $\lambda\in\mathbb{F}_{p^d}$. Hence, (3.10) has p^d solutions.

Case II, when $z=0$ and $y\neq 0$: We get that $y=tx$, where $t\in\mathbb{F}_{p^d}^*$, which implies that (3.10) has p^d-1 solutions.

Case III, when $yz\neq 0$: Since k is even, then $(\pi^{\frac{p^m-1}{4}})^{p^k}=\pi^{\frac{p^m-1}{4}}$. Together with $-1=\pi^{\frac{p^m-1}{2}}$ and by (3.10), we have

$$y^{p^k}-yx^{p^k-1}=\pm\pi^{\frac{p^m-1}{4}}(z^{p^k}-zx^{p^k-1}),$$

which is equivalent to

$$(y-\pi^{\frac{p^m-1}{4}}z)^{p^k}=(y-\pi^{\frac{p^m-1}{4}}z)x^{p^k-1} \tag{3.11}$$

or

$$(y+\pi^{\frac{p^m-1}{4}}z)^{p^k}=(y+\pi^{\frac{p^m-1}{4}}z)x^{p^k-1}. \tag{3.12}$$

Since $\gcd(p^k-1,p^m-1)=p^d-1$ and from (3.11) and (3.12), we get $y-\pi^{\frac{p^m-1}{4}}z=rx$ and $y+\pi^{\frac{p^m-1}{4}}z=r'x$, respectively, where $r,r'\in\mathbb{F}_{p^d}$. Hence, (3.11) and (3.12) have p^{m+d} solutions, respectively. Note that the common solutions of (3.11) and (3.12) are p^{2d}. Therefore, the total number of pairs $(y,z)\in\mathbb{F}_{p^d}^2$ satisfying (3.11) and (3.12) is $2p^{m+d}-p^{2d}$. In which the number satisfying $yz=0$ is $2p^d-1$. Then, when $yz\neq 0$, the total number of pairs $(y,z)\in\mathbb{F}_{p^d}^2$ satisfying (3.11) and (3.12) is $2p^{m+d}-p^{2d}-2p^d+1$.

As a result, we have

$$M_1=(p^d+p^d-1+2p^{m+d}-p^{2d}-2p^d+1)(p^m-1)=p^d(2p^m-p^d)(p^m-1).$$

Similarly, by (2.3),

$$\begin{aligned}
&\sum_{x,\alpha\in\mathbb{F}_{p^m}^*}T(\alpha,\psi(\alpha,x))T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\psi(\alpha,x))\\
=&\sum_{x,\alpha\in\mathbb{F}_{p^m}^*}\sum_{y,z\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\alpha(y^{p^k+1}+\pi^{\frac{p^k+1}{2}}z^{p^k+1})-\frac{1}{2}(\alpha x^{p^k-1}+\alpha^{p^{m-k}}x^{p^{m-k}-1})(y^2-\pi z^2))}\\
=&\sum_{x\in\mathbb{F}_{p^m}^*,y,z\in\mathbb{F}_{p^m}}\sum_{\alpha\in\mathbb{F}_{p^m}^*}\zeta_p^{\mathrm{Tr}_1^m(\frac{1}{2}\alpha x^{1-p^k}(-(x^{p^k-1}y-y^{p^k})^2+\pi(x^{p^k-1}z+\pi^{\frac{p^k-1}{2}}z^{p^k})^2))}\\
=&(p^m-1)M_2-(p^{2m}(p^m-1)-M_2)\\
=&p^mM_2-p^{3m}+p^{2m},
\end{aligned} \tag{3.13}$$

where

$$M_2=\#\{x\in\mathbb{F}_{p^m}^*,y,z\in\mathbb{F}_{p^m}\mid (x^{p^k-1}y-y^{p^k})^2=\pi(x^{p^k-1}z+\pi^{\frac{p^k-1}{2}}z^{p^k})^2\}.$$

For a given $x\in\mathbb{F}_{p^m}^*$, we study the following equation

$$(x^{p^k-1}y-y^{p^k})^2=\pi(x^{p^k-1}z+\pi^{\frac{p^k-1}{2}}z^{p^k})^2. \tag{3.14}$$

Case I, when $y=0$: In this case, we get that $z=0$ or $x^{p^k-1}=-\pi^{\frac{p^k-1}{2}}z^{p^k-1}$. Note that s is odd, then $p^d-1\nmid\frac{p^m-1}{2}$. Since $\frac{k}{d}$ is even, then $p^d-1\mid\frac{p^k-1}{2}$. Hence, there is no solution of $x^{p^k-1}=-\pi^{\frac{p^k-1}{2}}z^{p^k-1}$ for $z\in\mathbb{F}_{p^d}^*$. In this case, (3.14) has only one solution.

Case II, when $z=0$ and $y\neq 0$: We have that $x^{p^k-1}y-y^{p^k}=0$, which is equivalent to $x^{p^k-1}=y^{p^k-1}$, i.e., $y=tx$, where $t\in\mathbb{F}_{p^d}^*$. Hence, (3.14) has p^d-1 solutions.

Case III, when $yz\neq 0$: From case I, we get that $x^{p^k-1}+\pi^{\frac{p^k-1}{2}}z^{p^k-1}\neq 0$ for $z\neq 0$, which implies that $x^{p^k-1}y-y^{p^k}\neq 0$ by (3.14). Then there is no solution of (3.14).

Therefore, $M_2=p^d(p^m-1)$.

Similarly, by (2.3),

$$\begin{aligned}
&\sum_{x,\alpha\in\mathbb{F}_{p^m}^*}T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\psi(\alpha,x))^2\\
=&\sum_{x,\alpha\in\mathbb{F}_{p^m}^*}\sum_{y,z\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}(y^{p^k+1}+z^{p^k+1})+\frac{1}{2}\pi(\alpha x^{p^k-1}+\alpha^{p^{m-k}}x^{p^m-k-1})(y^2+z^2))}\\
=&\sum_{x\in\mathbb{F}_{p^m}^*,y,z\in\mathbb{F}_{p^m}}\sum_{\alpha\in\mathbb{F}_{p^m}^*}\zeta_p^{\mathrm{Tr}_1^m(\frac{1}{2}\pi\alpha x^{1-p^k}((x^{p^k-1}y+\pi^{\frac{p^k-1}{2}}y^{p^k})^2+(x^{p^k-1}z+\pi^{\frac{p^k-1}{2}}z^{p^k})^2))}\\
=&(p^m-1)M_3-(p^{2m}(p^m-1)-M_3)\\
=&p^mM_3-p^{3m}+p^{2m},
\end{aligned}\tag{3.15}$$

where

$$M_3=\#\{x\in\mathbb{F}_{p^m}^*,y,z\in\mathbb{F}_{p^m}\mid(x^{p^k-1}y+\pi^{\frac{p^k-1}{2}}y^{p^k})^2+(x^{p^k-1}z+\pi^{\frac{p^k-1}{2}}z^{p^k})^2=0\}.$$

For a given $x\in\mathbb{F}_{p^m}^*$, we investigate the following equation

$$(x^{p^k-1}y+\pi^{\frac{p^k-1}{2}}y^{p^k})^2=-(x^{p^k-1}z+\pi^{\frac{p^k-1}{2}}z^{p^k})^2,$$

which is equivalent to

$$-\pi^{\frac{p^k-1}{2}}(y-\pi^{\frac{p^m-1}{4}}z)^{p^k}=(y-\pi^{\frac{p^m-1}{4}}z)x^{p^k-1}\tag{3.16}$$

or

$$-\pi^{\frac{p^k-1}{2}}(y+\pi^{\frac{p^m-1}{4}}z)^{p^k}=(y+\pi^{\frac{p^m-1}{4}}z)x^{p^k-1}.\tag{3.17}$$

Case I, when $y-\pi^{\frac{p^m-1}{4}}z=0$ and $y+\pi^{\frac{p^m-1}{4}}z=0$: In this case, $y=z=0$. (3.16) has one solutions and (3.17) has one solution, respectively.

Case II, when $y-\pi^{\frac{p^m-1}{4}}z=0$ and $y+\pi^{\frac{p^m-1}{4}}z\neq 0$: (3.17) is equivalent to $-\pi^{\frac{p^k-1}{2}}(y+\pi^{\frac{p^m-1}{4}}z)^{p^k-1}=x^{p^k-1}$. Note that m/d is odd and k/d is even, then $p^d-1\nmid\frac{p^m-1}{2}$ and $p^d-1\mid\frac{p^k-1}{2}$. This implies that there is no solution of $-\pi^{\frac{p^k-1}{2}}(y+\pi^{\frac{p^m-1}{4}}z)^{p^k-1}=x^{p^k-1}$. Hence, (3.16) has p^m-1 solutions, (3.17) has no solution.

Case III, when $y-\pi^{\frac{p^m-1}{4}}z\neq 0$ and $y+\pi^{\frac{p^m-1}{4}}z=0$: We can discuss it by a similar way in the case II. Hence, (3.16) has no solution and (3.17) has p^m-1 solutions, respectively.

Case IV, when $(y-\pi^{\frac{p^m-1}{4}}z)(y+\pi^{\frac{p^m-1}{4}}z)\neq 0$: Similarly, we have that (3.16) and (3.17) have no solution, respectively.

So, we get that $M_3=(2p^m-1)(p^m-1)$.

From (3.8), (3.9), (3.13) and (3.15), we finish the proof. □

In the following, we define

$$N_{\varepsilon,\mu,i,j}=\{(\alpha,\beta)\mid(\alpha,\beta)\in N_{\varepsilon,i},(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)\in N_{\mu,j}\}\ \text{ and }\ n_{\varepsilon,\mu,i,j}=\mid N_{\varepsilon,\mu,i,j}\mid$$

for $\varepsilon,\mu=\pm1$, $0\leq i,j\leq 2$.

By Lemma 3.1, we have that if $ij\neq 0$, then $n_{\varepsilon,\mu,i,j}=0$. Moreover, it is easy to check that $n_{\varepsilon,\mu,i,j}=n_{\mu,\varepsilon,j,i}$ by the symmetry of $T(\alpha,\beta)$ and $T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)$. Hence, we need only to calculate $n_{\varepsilon,\mu,i,0}$. For convenience, we let $n_{\varepsilon,\mu,i,0}=n_{\varepsilon,\mu,i}$. With above preparation we can determine the value distribution of the exponential sum $S(\alpha,\beta)$.

Theorem 3.3. *With the notations given above. Then the value distribution of $S(\alpha,\beta)$ defined by (3.2) is as follows:*

value	*frequency*
$2p^m$	1
0	$\frac{1}{2}(p^{m+d}-3p^m+p^d+1)(p^m-1)/(p^d-1)$
$\pm(p^d-1)p^{\frac{m}{2}}$	$(p^{m-d}-1)(p^m-1)/(p^{2d}-1)$
$(\pm1-p^{\frac{d}{2}})p^{\frac{m}{2}}$	$\frac{1}{2}p^{\frac{m-d}{2}}(p^{\frac{m-d}{2}}-1)(p^m-1)$
$(\pm1+p^{\frac{d}{2}})p^{\frac{m}{2}}$	$\frac{1}{2}p^{\frac{m-d}{2}}(p^{\frac{m-d}{2}}+1)(p^m-1)$
$\pm2p^{\frac{m}{2}}$	$\frac{1}{4}(p^d-1)(p^{2m}-1)/(p^d+1)$

Proof. From Lemma 3.1 and Lemma 2.4, we get that

$$\begin{cases} n_{1,0}=n_{1,1,0}+n_{-1,1,0}+n_{1,1,1}+n_{-1,1,1}+n_{1,1,2}+n_{-1,1,2},\\ n_{-1,0}=n_{1,-1,0}+n_{-1,-1,0}+n_{1,-1,1}+n_{-1,-1,1}+n_{1,-1,2}+n_{-1,-1,2}.\end{cases}\tag{3.18}$$

By (3.2) and Lemma 3.1, we have that

$$\begin{cases} n_{1,1}=n_{1,1,1}+n_{1,-1,1},\\ n_{-1,1}=n_{-1,1,1}+n_{-1,-1,1},\\ n_{1,2}=n_{1,1,2}+n_{1,-1,2},\\ n_{-1,2}=n_{-1,1,2}+n_{-1,-1,2}.\end{cases}\tag{3.19}$$

Note that if the rank of $\mathrm{Tr}_d^m(\alpha x^{p^k+1}+\beta x^2)$ is odd and we compute the value of $T(\alpha,\beta)$ and $T(a\alpha,a\beta)$, where a is a nonsquare element in $\mathbb{F}_q$. By Lemma 2.2, we have

$$T(a\alpha,a\beta)=-T(\alpha,\beta),$$

and since s is odd, we have

$$\begin{cases} n_{1,1,0} = n_{-1,-1,0}, \\ n_{1,-1,0} = n_{-1,1,0}, \\ n_{1,1,1} = n_{1,-1,1}, \\ n_{-1,1,1} = n_{-1,-1,1}, \\ n_{1,1,2} = n_{-1,-1,2}, \\ n_{1,-1,2} = n_{-1,1,2}. \end{cases} \tag{3.20}$$

By Lemma 3.2 1), we get that

$$\begin{aligned} 4p^{3m} = \; & 4p^{2m} + 4p^m(n_{1,1,0} + n_{-1,-1,0}) + 2(p^{\frac{d}{2}}+1)^2 p^m(n_{1,1,1} + n_{-1,-1,1}) \\ & +2(p^{\frac{d}{2}}-1)^2 p^m(n_{1,-1,1} + n_{-1,1,1}) + 2(p^d+1)^2 p^m(n_{1,1,2} + n_{-1,-1,2}) \\ & +2(p^d-1)^2 p^m(n_{1,-1,2} + n_{-1,1,2}). \end{aligned} \tag{3.21}$$

By Lemma 3.2 2), we have that

$$\begin{aligned} & p^m(p^m-1)(2p^{m+d} - 2p^m + 2p^d - p^{2d} - 1) \\ = \; & (p^d-1)p^m((p^{\frac{d}{2}}+1)^2(n_{1,1,1} + n_{-1,-1,1}) + (p^{\frac{d}{2}}-1)^2(n_{1,-1,1} + n_{-1,1,1}) \\ & +(p^d+1)^3(n_{1,1,2} + n_{-1,-1,2}) + (p^d+1)(p^d-1)^2(n_{1,-1,2} + n_{-1,1,2})). \end{aligned} \tag{3.22}$$

Solving the system of equations consisting of (3.18)-(3.22), we have

$$n_{1,1,0} = n_{-1,-1,0} = \frac{1}{4}(p^d-1)(p^{2m}-1)/(p^d+1),$$

$$n_{1,-1,0} = n_{-1,1,0} = \frac{1}{4}(p^{m+d} - 3p^m + p^d + 1)(p^m-1)/(p^d-1),$$

$$n_{-1,1,1} = n_{-1,-1,1} = \frac{1}{4}p^{\frac{m-d}{2}}(p^{\frac{m-d}{2}}-1)(p^m-1),$$

$$n_{1,1,1} = n_{1,-1,1} = \frac{1}{4}p^{\frac{m-d}{2}}(p^{\frac{m-d}{2}}+1)(p^m-1),$$

$$n_{1,-1,2} = n_{-1,1,2} = \frac{1}{2}(p^{m-d}-1)(p^m-1)/(p^{2d}-1),$$

$$n_{1,1,2} = n_{-1,-1,2} = 0.$$

We complete the proof. □

Therefore, we can give the weight distribution of cyclic code $\mathcal{C}$.

Theorem 3.4. *With the notations given above. Then $\mathcal{C}$ defined by (1.1) is a cyclic code over $\mathbb{F}_p$ with length p^m-1 and dimension $2m$. Moreover, the weight distribution of cyclic code $\mathcal{C}$ is given in Table 1.*

Proof. By (3.1), we have

$$wt(\mathbf{c}(\alpha,\beta)) = p^m - p^{m-1} - \frac{1}{2p}\sum_{u\in\mathbb{F}_p^*} S(u\alpha, u\beta).$$

Table 1: weight distribution of $\mathcal{C}$ for $1 \le v_2(m) < v_2(k)$

Weight	Frequency
0	1
$p^{m-1}(p-1)+\frac{1}{2}(p-1)(p^d-1)p^{\frac{m}{2}-1}$	$(p^{m-d}-1)(p^m-1)/(p^{2d}-1)$
$p^{m-1}(p-1)-\frac{1}{2}(p-1)(p^d-1)p^{\frac{m}{2}-1}$	$(p^{m-d}-1)(p^m-1)/(p^{2d}-1)$
$p^{m-1}(p-1)+\frac{1}{2}(p-1)(p^{\frac{d}{2}}-1)p^{\frac{m}{2}-1}$	$\frac{1}{2}p^{\frac{m-d}{2}}(p^{\frac{m-d}{2}}-1)(p^m-1)$
$p^{m-1}(p-1)+\frac{1}{2}(p-1)(p^{\frac{d}{2}}+1)p^{\frac{m}{2}-1}$	$\frac{1}{2}p^{\frac{m-d}{2}}(p^{\frac{m-d}{2}}-1)(p^m-1)$
$p^{m-1}(p-1)-\frac{1}{2}(p-1)(p^{\frac{d}{2}}+1)p^{\frac{m}{2}-1}$	$\frac{1}{2}p^{\frac{m-d}{2}}(p^{\frac{m-d}{2}}+1)(p^m-1)$
$p^{m-1}(p-1)-\frac{1}{2}(p-1)(p^{\frac{d}{2}}-1)p^{\frac{m}{2}-1}$	$\frac{1}{2}p^{\frac{m-d}{2}}(p^{\frac{m-d}{2}}+1)(p^m-1)$
$p^{m-1}(p-1)+p^{\frac{m}{2}-1}(p-1)$	$\frac{1}{4}(p^d-1)(p^{2m}-1)/(p^d+1)$
$p^{m-1}(p-1)-p^{\frac{m}{2}-1}(p-1)$	$\frac{1}{4}(p^d-1)(p^{2m}-1)/(p^d+1)$
$p^{m-1}(p-1)$	$\frac{1}{2}(p^{m+d}-3p^m+p^d+1)(p^m-1)/(p^d-1)$

Note that m is even, then $u_p = \pi^{\frac{p^m-1}{p-1}} = \pi^{p^{m-1}+\cdots+1}$ is a square element in $\mathbb{F}_{p^m}$. For a given $u \in \mathbb{F}_p^*$, we have that $u \in SQ$. Since k is even, then $u^{\frac{p^k+1}{2}} = u$. So we have

$$\begin{aligned}
S(u\alpha, u\beta) &= \sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{\mathrm{Tr}_1^m(u\alpha x^{p^k+1}+u\beta x^2)} + \zeta_p^{\mathrm{Tr}_1^m(u\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-u\beta\pi x^2)}\right)\\
&= \sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{\mathrm{Tr}_1^m(\alpha(u^{\frac{1}{2}}x)^{p^k+1}+\beta(u^{\frac{1}{2}}x)^2)} + \zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}(u^{\frac{1}{2}}x)^{p^k+1}-\beta\pi(u^{\frac{1}{2}}x)^2)}\right)\\
&= \sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1}+\beta x^2)} + \zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-\beta\pi x^2)}\right)\\
&= S(\alpha,\beta).
\end{aligned}$$

Therefore, we obtain that

$$wt(\mathbf{c}(\alpha,\beta)) = p^m - p^{m-1} - \frac{p-1}{2p}S(\alpha,\beta).$$

By Theorem 3.3, we get the result. □

In the following, we give an example to verify our result in Theorem 3.4.

Example 3.5. *Let $p=3$, $m=6$, $k=4$, the code $\mathcal{C}$ is a $[728, 12, 414]$ cyclic code over $\mathbb{F}_3$ with weight enumerator*

$$1 + 728X^{414} + 32760X^{450} + 139048X^{468} + 199472X^{486} + 132496X^{504} + 26208X^{522} + 728X^{558},$$

which confirms the weight distribution in Table 1.

3.2 The weight distribution of $\mathcal{C}$ for $v_2(k) < v_2(m)$

In this subsection, we always assume that $v_2(k) < v_2(m)$. To determine the value distribution of $S(\alpha,\beta)$, we need some identities on $S(\alpha,\beta)$.

Lemma 3.6. *With the notations given above, we have the following result:*

1. $\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta) = 2p^{2m}$.

2. $\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta)^2 = \begin{cases} 4p^{3m}, & \text{if } p^k \equiv 1 \pmod 4; \\ 4p^{2m}, & \text{if } p^k \equiv 3 \pmod 4. \end{cases}$

3. $\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta)^3 = \begin{cases} 2p^{2m}(7p^m-3) + 2p^{2m+d}(p^m-1), & \text{if } p^k \equiv 1 \pmod 4; \\ 2p^{2m}(p^m+3) + 2p^{2m+d}(p^m-1), & \text{if } p^k \equiv 3 \pmod 4. \end{cases}$

4. $(p^d-1)\sum_{(\alpha,\beta)\in N_1} S(\alpha,\beta) + (p^{2d}-1)\sum_{(\alpha,\beta)\in N_2} S(\alpha,\beta) = p^m(p^d-1)(p^m-1)$.

Proof. 1. We compute

$$\begin{aligned} \sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta) &= \sum_{\alpha,\beta\in\mathbb{F}_{p^m}} \left(T(\alpha,\beta) + T(\pi^{\frac{p^k+1}{2}}\alpha, -\pi\beta) \right) \\ &= \sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta) + \sum_{\alpha',\beta'\in\mathbb{F}_{p^m}} T(\alpha',\beta') \\ &= 2\sum_{x\in\mathbb{F}_{p^m}} \sum_{\alpha\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1})} \sum_{\beta\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\beta x^2)} \\ &= 2p^{2m}. \end{aligned}$$

2. By (3.2), we get that

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta)^2 = \sum_{\alpha,\beta\in\mathbb{F}_{p^m}} \left(T(\alpha,\beta)^2 + 2T(\alpha,\beta)T(\pi^{\frac{p^k+1}{2}}\alpha, -\pi\beta) + T(\pi^{\frac{p^k+1}{2}}\alpha, -\pi\beta)^2 \right). \quad (3.23)$$

On one hand,

$$\begin{aligned} \sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)^2 &= \sum_{x,y\in\mathbb{F}_{p^m}} \sum_{\alpha\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha(x^{p^k+1}+y^{p^k+1}))} \sum_{\beta\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha(x^2+y^2))} \\ &= Mp^{2m}, \end{aligned} \quad (3.24)$$

where $M = \#\{(x,y)\in\mathbb{F}_{p^m}^2 \mid x^2+y^2=0, x^{p^k+1}+y^{p^k+1}=0\}$. Together with Lemma 2.5 ii) and (3.24), we obtain that

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)^2 = \begin{cases} p^{2m}(2p^m-1), & \text{if } p^k \equiv 1 \pmod 4; \\ p^{2m}, & \text{if } p^k \equiv 3 \pmod 4. \end{cases}$$

Similarly,

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\pi^{\frac{p^k+1}{2}}\alpha, -\pi\beta)^2 = \begin{cases} p^{2m}(2p^m-1), & \text{if } p^k \equiv 1 \pmod 4; \\ p^{2m}, & \text{if } p^k \equiv 3 \pmod 4. \end{cases}$$

On the other hand,

$$\begin{aligned}
&\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)\\
=\;& \sum_{x,y\in\mathbb{F}_{p^m}}\sum_{\alpha\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha(x^{p^k+1}+\pi^{\frac{p^k+1}{2}}y^{p^k+1}))}\sum_{\beta\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\beta(x^2-\pi y^2))}\\
=\;& \#\{(x,y)\in\mathbb{F}_{p^m}^2 \mid x^2-\pi y^2=0, x^{p^k+1}+\pi^{\frac{p^k+1}{2}}y^{p^k+1}=0\}\cdot p^{2m}\\
=\;& \#\{(x,y)\in\mathbb{F}_{p^m}^2 \mid x^2-\pi y^2=0, 2\pi^{\frac{p^k+1}{2}}y^{p^k+1}=0\}\cdot p^{2m}\\
=\;& p^{2m}.
\end{aligned}$$

Hence, from (3.23), we get

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta)^2=\begin{cases} 4p^{3m}, & \text{if } p^k\equiv 1 \pmod 4;\\ 4p^{2m}, & \text{if } p^k\equiv 3 \pmod 4.\end{cases}$$

3. By (3.2),

$$\begin{aligned}
\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta)^3 \;=\;& \sum_{\alpha,\beta\in\mathbb{F}_{p^m}} (T(\alpha,\beta)^3+3T(\alpha,\beta)^2T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)\\
&+3T(\alpha,\beta)T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)^2+T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)^3). \qquad (3.25)
\end{aligned}$$

It is easy to check that

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)^3=\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)^3 \qquad (3.26)$$

and

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)^2T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)=\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)^2. \qquad (3.27)$$

By Lemma 2.4 ii),

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)^3=p^{3m}+p^{2m+d}(p^m-1). \qquad (3.28)$$

By Lemma 2.6, we have

$$\begin{aligned}
&\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} T(\alpha,\beta)^2T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\beta)\\
=\;& \sum_{x,y,z\in\mathbb{F}_{p^m}}\sum_{\alpha\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha(x^{p^k+1}+y^{p^k+1}+\pi^{\frac{p^k+1}{2}}z^{p^k+1}))}\sum_{\beta\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\beta(x^2+y^2-\pi z^2))}\\
=\;& \#\{(x,y,z)\in\mathbb{F}_{p^m}^3 \mid x^2+y^2-\pi z^2=0, x^{p^k+1}+y^{p^k+1}+\pi^{\frac{p^k+1}{2}}z^{p^k+1})=0\}\cdot p^{2m}\\
=\;& \begin{cases} (2p^m-1)p^{2m}, & \text{if } p^k\equiv 1 \pmod 4;\\ p^{2m}, & \text{if } p^k\equiv 3 \pmod 4.\end{cases} \qquad (3.29)
\end{aligned}$$

Combining (3.25)-(3.29), we get

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta)^3 = \begin{cases} 2p^{2m}(7p^m-3)+2p^{2m+d}(p^m-1), & \text{if } p^k\equiv 1 \pmod 4; \\ 2p^{2m}(p^m+3)+2p^{2m+d}(p^m-1), & \text{if } p^k\equiv 3 \pmod 4. \end{cases}$$

4. By (2.5) and (3.2),

$$\begin{aligned} & (p^d-1)\sum_{(\alpha,\beta)\in N_1} S(\alpha,\beta)+(p^{2d}-1)\sum_{(\alpha,\beta)\in N_2} S(\alpha,\beta) \\ = & \sum_{x,\alpha\in\mathbb{F}_{p^m}^*}\left(T(\alpha,\psi(\alpha,x))+T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\psi(\alpha,x))\right). \end{aligned} \tag{3.30}$$

By (2.4),

$$\begin{aligned} \sum_{x,\alpha\in\mathbb{F}_{p^m}^*} T(\alpha,\psi(\alpha,x)) &= \sum_{x,\alpha\in\mathbb{F}_{p^m}^*}\sum_{y\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha y^{p^k+1}-\frac{1}{2}x^{-1}(\alpha x^{p^k}+\alpha^{p^{m-k}}x^{p^{m-k}})y^2)} \\ &= \sum_{x,\alpha\in\mathbb{F}_{p^m}^*}\sum_{y\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\alpha y^{p^k+1}-\frac{1}{2}x^{-1}\alpha x^{p^k}y^2-\frac{1}{2}x^{-p^k}\alpha x y^{2p^k})} \\ &= \sum_{x\in\mathbb{F}_{p^m}^*,y\in\mathbb{F}_{p^m}}\sum_{\alpha\in\mathbb{F}_{p^m}^*} \zeta_p^{\mathrm{Tr}_1^m(-\frac{1}{2}\alpha(y^2x^{1-p^k}(x^{p^k-1}-y^{p^k-1})^2))} \\ &= (p^m-1)M-(p^m(p^m-1)-M) \\ &= p^mM-p^{2m}+p^m, \end{aligned} \tag{3.31}$$

where

$$\begin{aligned} M &= \#\{x\in\mathbb{F}_{p^m}^*, y\in\mathbb{F}_{p^m} \mid y^2(x^{p^k-1}-y^{p^k-1})^2=0\} \\ &= \#\{x\in\mathbb{F}_{p^m}^*, y\in\mathbb{F}_{p^m} \mid \ y=\lambda x, \text{where } \lambda\in\mathbb{F}_{p^d}\} \\ &= (p^m-1)p^d. \end{aligned} \tag{3.32}$$

On the other hand, by (2.5),

$$\begin{aligned} \sum_{x,\alpha\in\mathbb{F}_{p^m}^*} T(\pi^{\frac{p^k+1}{2}}\alpha,-\pi\psi(\alpha,x)) &= \sum_{x,\alpha\in\mathbb{F}_{p^m}^*}\sum_{y\in\mathbb{F}_{p^m}} \zeta_p^{\mathrm{Tr}_1^m(\pi^{\frac{p^k+1}{2}}\alpha y^{p^k+1}+\pi\frac{1}{2}x^{-1}(\alpha x^{p^k}+\alpha^{p^{m-k}}x^{p^{m-k}})y^2)} \\ &= \sum_{x\in\mathbb{F}_{p^m}^*,y\in\mathbb{F}_{p^m}}\sum_{\alpha\in\mathbb{F}_{p^m}^*} \zeta_p^{\mathrm{Tr}_1^m(\frac{1}{2}\pi\alpha(y^2x^{1-p^k}(x^{p^k-1}+\pi^{\frac{p^k-1}{2}}y^{p^k-1})^2))} \\ &= (p^m-1)I-(p^m(p^m-1)-I) \\ &= p^mI-p^{2m}+p^m, \end{aligned} \tag{3.33}$$

where

$$I = \#\{x\in\mathbb{F}_{p^m}^*, y\in\mathbb{F}_{p^m} \mid y^2(x^{p^k-1}+\pi^{\frac{p^k-1}{2}}y^{p^k-1})^2=0\}$$

$$= \quad \#\{x \in \mathbb{F}_{p^m}^*, y \in \mathbb{F}_{p^m} \mid y = 0 \text{ or } (\frac{x}{y})^{p^k-1} = -\pi^{\frac{p^k-1}{2}}\}.$$

Since $\frac{k}{d}$ is odd and m is even, then $p^d - 1 \nmid \frac{p^k-1}{2} + \frac{p^m-1}{2}$, which implies that there is no solution of $(\frac{x}{y})^{p^k-1} = -\pi^{\frac{p^k-1}{2}}$. Hence, $I = p^m - 1$. Together with (3.30), (3.31) and (3.33), we have that

$$(p^d-1)\sum_{(\alpha,\beta)\in N_1} S(\alpha,\beta) + (p^{2d}-1)\sum_{(\alpha,\beta)\in N_2} S(\alpha,\beta) = p^m(p^d-1)(p^m-1).$$

□

Theorem 3.7. *With the notations given above. Then the value distribution of $S(\alpha,\beta)$ defined by (3.2) is as follows:*

value	*frequency*
$2p^m$	1
0	$\frac{1}{2}(p^{m+d} - 3p^m + p^d + 1)(p^m-1)/(p^d-1)$
$\pm(p^d-1)p^{\frac{m}{2}}$	$(p^{\frac{m}{2}-d} \pm 1)(p^{\frac{m}{2}} \mp 1)(p^m-1)/(p^{2d}-1)$
$(\pm\sqrt{q^*}-1)p^{\frac{m}{2}}$	$\frac{1}{2}p^{\frac{m}{2}-d}(p^{\frac{m}{2}}-1)(p^m-1)$
$(\pm\sqrt{q^*}+1)p^{\frac{m}{2}}$	$\frac{1}{2}p^{\frac{m}{2}-d}(p^{\frac{m}{2}}+1)(p^m-1)$
$\pm 2p^{\frac{m}{2}}$	$\frac{1}{4}(p^{\frac{m}{2}} \pm 1)^2(p^d-1)(p^m-1)/(p^d+1)$

Proof. From Lemma 3.1 and Lemma 2.4, we get that

$$\begin{cases} n_{1,0} = n_{1,1,0} + n_{-1,1,0} + n_{1,1,1} + n_{-1,1,1} + n_{1,1,2} + n_{-1,1,2}, \\ n_{-1,0} = n_{1,-1,0} + n_{-1,-1,0} + n_{1,-1,1} + n_{-1,-1,1} + n_{1,-1,2} + n_{-1,-1,2}. \end{cases} \tag{3.34}$$

By (3.2) and Lemma 3.1,

$$\begin{cases} n_{1,1} = n_{1,1,1} + n_{1,-1,1}, \\ n_{-1,1} = n_{-1,1,1} + n_{-1,-1,1}, \\ n_{1,2} = n_{1,1,2} + n_{1,-1,2}, \\ n_{-1,2} = n_{-1,1,2} + n_{-1,-1,2}. \end{cases} \tag{3.35}$$

Note that if the rank of $\mathrm{Tr}_d^m(\alpha x^{p^k+1} + \beta x^2)$ is odd and we compute the value of $T(\alpha,\beta)$ and $T(a\alpha, a\beta)$, where a is a nonsquare element in $\mathbb{F}_q$. Then, by Lemma 2.2, we have

$$T(a\alpha, a\beta) = -T(\alpha,\beta),$$

and since s is even , we get

$$\begin{cases} n_{1,1,1} = n_{-1,1,1}, \\ n_{1,-1,1} = n_{-1,-1,1}. \end{cases} \tag{3.36}$$

On the other hand, we have

$$\begin{aligned} \sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta) = 2p^m + 2p^{\frac{m}{2}}(n_{1,1,0} - n_{-1,-1,0} + 2(\sqrt{q^*}+1)(n_{1,1,1} - n_{-1,-1,1}) + \\ 2(\sqrt{q^*}-1)(n_{1,-1,1} - n_{-1,1,1}) + (p^d+1)(n_{1,1,2} - n_{-1,-1,2}) + (p^d-1)(n_{1,-1,2} - n_{-1,1,2})), \end{aligned} \tag{3.37}$$

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta)^2 = 4p^{2m} + 2p^m(2(n_{1,1,0}+n_{-1,-1,0}) + (\sqrt{q^*}+1)^2(n_{1,1,1}+n_{-1,-1,1}) + (\sqrt{q^*}-1)^2(n_{1,-1,1}-n_{-1,1,1}) + (p^d+1)^2(n_{1,1,2}+n_{-1,-1,2}) + (p^d-1)^2(n_{1,-1,2}+n_{-1,1,2})), \tag{3.38}$$

$$\sum_{\alpha,\beta\in\mathbb{F}_{p^m}} S(\alpha,\beta)^3 = 8p^{3m} + 2p^{\frac{3m}{2}}(4(n_{1,1,0}-n_{-1,-1,0}) + (\sqrt{q^*}+1)^3(n_{1,1,1}-n_{-1,-1,1}) + (\sqrt{q^*}-1)^3(n_{1,-1,1}-n_{-1,1,1}) + (p^d+1)^3(n_{1,1,2}-n_{-1,-1,2}) + (p^d-1)^3(n_{1,-1,2}-n_{-1,1,2})), \tag{3.39}$$

$$(p^d-1)\sum_{(\alpha,\beta)\in N_1} S(\alpha,\beta) + (p^{2d}-1)\sum_{(\alpha,\beta)\in N_2} S(\alpha,\beta) = (p^d-1)p^{\frac{m}{2}}((\sqrt{q^*}+1)(n_{1,1,1}-n_{-1,-1,1}) + (\sqrt{q^*}-1)(n_{1,-1,1}-n_{-1,1,1}) + (p^d+1)^2(n_{1,1,2}-n_{-1,-1,2}) + (p^{2d}-1)(n_{1,-1,2}-n_{-1,1,2})). \tag{3.40}$$

Applying Lemma 3.6 $1)-4)$ and solving the system equations consisting of (3.34)-(3.40), we get

$$n_{1,1,0} = \frac{1}{4}(p^{\frac{m}{2}}+1)^2(p^d-1)(p^m-1)/(p^d+1),$$

$$n_{-1,-1,0} = \frac{1}{4}(p^{\frac{m}{2}}-1)^2(p^d-1)(p^m-1)/(p^d+1),$$

$$n_{1,-1,0} = n_{-1,1,0} = \frac{1}{4}(p^{m+d}-3p^m+p^d+1)(p^m-1)/(p^d-1),$$

$$n_{1,1,1} = n_{-1,1,1} = \frac{1}{4}p^{\frac{m}{2}-d}(p^{\frac{m}{2}}+1)(p^m-1),$$

$$n_{1,-1,1} = n_{-1,-1,1} = \frac{1}{4}p^{\frac{m}{2}-d}(p^{\frac{m}{2}}-1)(p^m-1),$$

$$n_{1,-1,2} = \frac{1}{2}(p^{\frac{m}{2}-d}+1)(p^{\frac{m}{2}}-1)(p^m-1)/(p^{2d}-1),$$

$$n_{-1,1,2} = \frac{1}{2}(p^{\frac{m}{2}-d}-1)(p^{\frac{m}{2}}+1)(p^m-1)/(p^{2d}-1),$$

$$n_{1,1,2} = n_{-1,-1,2} = 0.$$

This finishes the proof. □

Therefore, we can determine the weight distribution of cyclic code $\mathcal{C}$.

Theorem 3.8. *With the notations given above.*

i) If k is odd, then $\mathcal{C}$ defined by (1.1) is a cyclic code over $\mathbb{F}_p$ with length p^m-1 and dimension $2m$. Moreover, the weight distribution of cyclic code $\mathcal{C}$ is given in Table 2.

ii) If k is even, then $\mathcal{C}$ defined by (1.1) is a cyclic code over $\mathbb{F}_p$ with length p^m-1 and dimension $2m$. Moreover, the weight distribution of cyclic code $\mathcal{C}$ is given in Table 3.

Proof. By (3.1), we have

$$wt(\mathbf{c}(\alpha,\beta)) = p^m - p^{m-1} - \frac{1}{2p}\sum_{u\in\mathbb{F}_p^*} S(u\alpha,u\beta).$$

Table 2: weight distribution of $\mathcal{C}$ for odd k

value	frequency
0	1
$p^m - p^{m-1}$	$\frac{1}{2}(p^{m+d} - 3p^m + p^d + 1)(p^m - 1)/(p^d - 1)$
$p^{m-1}(p-1) - \frac{1}{2}(p-1)(p^d-1)p^{\frac{m}{2}-1}$	$(p^{\frac{m}{2}-d} + 1)(p^{\frac{m}{2}} - 1)(p^m - 1)/(p^{2d} - 1)$
$p^{m-1}(p-1) + \frac{1}{2}(p-1)(p^d-1)p^{\frac{m}{2}-1}$	$(p^{\frac{m}{2}-d} - 1)(p^{\frac{m}{2}} + 1)(p^m - 1)/(p^{2d} - 1)$
$p^{m-1}(p-1) + \frac{1}{2}(p-1)p^{\frac{m}{2}-1}$	$p^{\frac{m}{2}-d}(p^{\frac{m}{2}} - 1)(p^m - 1)$
$p^{m-1}(p-1) - \frac{1}{2}(p-1)p^{\frac{m}{2}-1}$	$p^{\frac{m}{2}-d}(p^{\frac{m}{2}} + 1)(p^m - 1)$
$p^{m-1}(p-1) + (p-1)p^{\frac{m}{2}-1}$	$\frac{1}{4}(p^{\frac{m}{2}} - 1)^2(p^d - 1)(p^m - 1)/(p^d + 1)$
$p^{m-1}(p-1) - (p-1)p^{\frac{m}{2}-1}$	$\frac{1}{4}(p^{\frac{m}{2}} + 1)^2(p^d - 1)(p^m - 1)/(p^d + 1)$

Table 3: weight distribution of $\mathcal{C}$ for even k

Weight	Frequency
0	1
$p^m - p^{m-1}$	$\frac{1}{2}(p^{m+d} - 3p^m + p^d + 1)(p^m - 1)/(p^d - 1)$
$p^{m-1}(p-1) + \frac{1}{2}(p-1)(p^d-1)p^{\frac{m}{2}-1}$	$(p^{\frac{m}{2}-d} - 1)(p^{\frac{m}{2}} + 1)(p^m - 1)/(p^{2d} - 1)$
$p^{m-1}(p-1) - \frac{1}{2}(p-1)(p^d-1)p^{\frac{m}{2}-1}$	$(p^{\frac{m}{2}-d} + 1)(p^{\frac{m}{2}} - 1)(p^m - 1)/(p^{2d} - 1)$
$p^{m-1}(p-1) - \frac{1}{2}(p-1)(p^{\frac{d}{2}}-1)p^{\frac{m}{2}-1}$	$\frac{1}{2}p^{\frac{m}{2}-d}(p^{\frac{m}{2}} - 1)(p^m - 1)$
$p^{m-1}(p-1) + \frac{1}{2}(p-1)(p^{\frac{d}{2}}+1)p^{\frac{m}{2}-1}$	$\frac{1}{2}p^{\frac{m}{2}-d}(p^{\frac{m}{2}} - 1)(p^m - 1)$
$p^{m-1}(p-1) - \frac{1}{2}(p-1)(p^{\frac{d}{2}}+1)p^{\frac{m}{2}-1}$	$\frac{1}{2}p^{\frac{m}{2}-d}(p^{\frac{m}{2}} + 1)(p^m - 1)$
$p^{m-1}(p-1) + \frac{1}{2}(p-1)(p^{\frac{d}{2}}-1)p^{\frac{m}{2}-1}$	$\frac{1}{2}p^{\frac{m}{2}-d}(p^{\frac{m}{2}} + 1)(p^m - 1)$
$p^{m-1}(p-1) + (p-1)p^{\frac{m}{2}-1}$	$\frac{1}{4}(p^{\frac{m}{2}} - 1)^2(p^d - 1)(p^m - 1)/(p^d + 1)$
$p^{m-1}(p-1) - (p-1)p^{\frac{m}{2}-1}$	$\frac{1}{4}(p^{\frac{m}{2}} + 1)^2(p^d - 1)(p^m - 1)/(p^d + 1)$

If k is odd, then d is odd and $u_p = \pi^{\frac{p^m-1}{p-1}} = \pi^{\frac{p^m-1}{p^d-1}\cdot\frac{p^d-1}{p-1}}$ is a nonsquare element in $\mathbb{F}_{p^d}$. So we have $\eta_d(u_p) = -1$. If $u \in SQ_p$, then $u^{\frac{p^k+1}{2}} = u$. Hence, by Lemma 2.2, we have

$$\begin{aligned}
&\sum_{u\in\mathbb{F}_p^*} S(u\alpha, u\beta)\\
=\ &\sum_{u\in\mathbb{F}_p^*}\sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{\mathrm{Tr}_1^m(u\alpha x^{p^k+1}+u\beta x^2)} + \zeta_p^{\mathrm{Tr}_1^m(u\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-u\beta\pi x^2)}\right)\\
=\ &\sum_{u\in SQ_p}\sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{\mathrm{Tr}_1^m(\alpha(u^{\frac{1}{2}}x)^{p^k+1}+\beta(u^{\frac{1}{2}}x)^2)} + \zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}(u^{\frac{1}{2}}x)^{p^k+1}-\beta\pi(u^{\frac{1}{2}}x)^2)}\right)\\
+\ &\sum_{u\in SQ_p}\sum_{x\in\mathbb{F}_{p^m}} \left(\zeta_p^{u_p\mathrm{Tr}_1^m(\alpha(u^{\frac{1}{2}}x)^{p^k+1}+\beta(u^{\frac{1}{2}}x)^2)} + \zeta_p^{u_p\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}(u^{\frac{1}{2}}x)^{p^k+1}-\beta\pi(u^{\frac{1}{2}}x)^2)}\right)
\end{aligned}$$

$$
\begin{aligned}
&= \sum_{u\in SQ_p}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1}+\beta x^2)}+\zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-\beta\pi x^2)}\right)\\
&+ \sum_{u\in SQ_p}\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{u_p\mathrm{Tr}_1^m(\alpha x^{p^k+1}+\beta x^2)}+\zeta_p^{u_p\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-\beta\pi x^2)}\right)\\
&= \frac{p-1}{2}\left((1+\eta_d(u_p)^r\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1}+\beta x^2)}+(1+\eta_d(u_p)^{r'}\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-\beta\pi x^2)}\right),
\end{aligned}
$$

where r and r' are the rank of $\mathrm{Tr}_d^m(\alpha x^{p^k+1}+\beta x^2)$ and $\mathrm{Tr}_d^m(\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-\beta\pi x^2)$, respectively. Note that $\eta_d(u_p)=-1$ and s is even. By Lemma 3.1, we have if $r=s-1$, then $r'=s$ and

$$
\sum_{u\in\mathbb{F}_p^*}S(u\alpha,u\beta)=(p-1)\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-\beta\pi x^2)}.
$$

If $r'=s-1$, then $r=s$ and

$$
\sum_{u\in\mathbb{F}_p^*}S(u\alpha,u\beta)=(p-1)\sum_{x\in\mathbb{F}_{p^m}}\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1}+\beta x^2)}.
$$

Otherwise,

$$
\sum_{u\in\mathbb{F}_p^*}S(u\alpha,u\beta)=(p-1)\sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1}+\beta x^2)}+\zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-\beta\pi x^2)}\right).
$$

Therefore, by Theorem 3.7, we obtain the weight distribution of cyclic code $\mathcal{C}$ for odd k.

If k is even, then $u^{\frac{p^k+1}{2}}=u$ for all $u\in\mathbb{F}_p$. Since m is even, then u is a square element in $\mathbb{F}_{p^m}$. Hence, we have

$$
\begin{aligned}
S(u\alpha,u\beta) &= \sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m(u\alpha x^{p^k+1}+u\beta x^2)}+\zeta_p^{\mathrm{Tr}_1^m(u\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-u\beta\pi x^2)}\right)\\
&= \sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m(\alpha(u^{\frac12}x)^{p^k+1}+\beta(u^{\frac12}x)^2)}+\zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}(u^{\frac12}x)^{p^k+1}-\beta\pi(u^{\frac12}x)^2)}\right)\\
&= \sum_{x\in\mathbb{F}_{p^m}}\left(\zeta_p^{\mathrm{Tr}_1^m(\alpha x^{p^k+1}+\beta x^2)}+\zeta_p^{\mathrm{Tr}_1^m(\alpha\pi^{\frac{p^k+1}{2}}x^{p^k+1}-\beta\pi x^2)}\right)=S(\alpha,\beta).
\end{aligned}
$$

Therefore,

$$
wt(\mathbf{c}(\alpha,\beta))=p^m-p^{m-1}-\frac{p-1}{2p}S(\alpha,\beta).
$$

By Theorem 3.7, we get the weight distribution of cyclic code $\mathcal{C}$ for even k. □

In the following, we give an example to verify the result in Theorem 3.8 for the case of odd k. For the case of even k, we are not able to give an example to verify the result in Theorem 3.8 because of our limited computation ability.

Example 3.9. *Let $p = 3$, $m = 6$, $k = 1$, the code $\mathcal{C}$ is a $[728, 12, 468]$ cyclic code over $\mathbb{F}_3$ with weight enumerator*

$$1 + 95004X^{468} + 183456X^{477} + 728X^{486} + 170352X^{495} + 81900X^{504},$$

which confirms the weight distribution in Table 2.

References

[1] P. Delsarte, "On subfield subcodes of modified Reed-Solomon codes," *IEEE Trans. Inform. Theory*, **21**(5), 575-576(1975).

[2] C. Ding, Y. Liu, C. Ma, L. Zeng, "The weight distributions of the duals of cyclic codes with two zeros," *IEEE Trans. Inform. Theory*, **57**(12), 8000-8006(2011).

[3] C. Ding, J. Yang, "Hamming weight in irrecducible codes," *Discret. Math.* **313**(4), 434-446(2013).

[4] S. Draper, X. Hou, "explicit evaluation of certain exponential sums of quadratic functions over $\mathbb{F}_{p^m}$, p odd," http://arxiv.org/pdf/0708.3619v1.pdf.

[5] K. Feng, J. Luo, "Weight distribution of some reducible cyclic codes," *Finite Fields Appl.* **14**(2), 390-409(2008).

[6] T. Feng, "On cyclic codes of length $2^{2^r} - 1$ with two zeros whose dual codes have three weights," *Des. Codes Cryptogr.* **62**(3), 253-258(2012).

[7] T. Helleseth, A. Kholosha, "Monomial and quadratic bent functions over the finite fields of odd characteristic", *IEEE Trans. Inform. Theory,* **52**(5), 2018-2032(2006).

[8] R. Lidl, H. Niederreiter, "Finite Fields," Ency clopedia of Mathematics, **20**, Cambridge University Press, Cambridge(1983).

[9] Y. Liu, H. Yan, C. Liu, "A class of six-weight cyclic codes and their weight distribution," http://arxiv.org/pdf/1311.3391v2.pdf.

[10] J. Luo, K. Feng, "Cyclic codes and sequences from generalized Coulter-Matthews function," *IEEE Trans. Inform. Theory*, **54**(12), 5345-5353(2008).

[11] J. Luo, K. Feng, "On the weight distributions of two classes of cyclic codes," *IEEE Trans. Inform. Theory*, **54**(12), 5332-5344(2008).

[12] C. Ma, L. zeng, Y. Liu, D. Feng, C. Ding, "The weight enumerator of a class of cyclic codes," *IEEE Trans. Inform. Theory*, **57**(1), 397-402(2011).

[13] F. MacWilliams, N. Sloane, "The Theory of Error-Correcting Codes," North-Holland Publishing, Amsterdam(1997).

[14] A. Rao, N. Pinnawala, "A family of two-weight irreducible cyclic codes," *IEEE Trans. Inform. Theory*, **56**(6), 2568-2570(2010).

[15] G. Vega, "The weight distribution of an extended class of reducible cyclic codes," *IEEE Trans. Inform. Theory*, **58**(7), 4862-4869(2012).

[16] G. Vega, J. Wolfmann, "New classes of 2-weight cyclic codes," *Des. Codes Cryptogr.* **42**(3), 327-334(2007).

[17] B. Wang, C. Tang, Y. Qi , Y. Yang , M. Xu, "The weight distributions of cyclic codes and elliptic curves," *IEEE Trans. Inform. Theory*, **58**(12), 7253-7259(2012).

[18] M. Xiong, "The weight distributions of a class of cyclic codes II," *Des. Codes Cryptogr.* (2012). doi: 10.1007/s10623-012-9785-0.

[19] M. Xiong, "The weight distributions of a class of cyclic codes III," *Finite Fields Appl.* **21**, 84-96(2013).

[20] M. Xiong, "The weight distributions of a class of cyclic codes," *Finite Fields Appl.* **18**(5), 933-945(2012).

[21] L. Yu, H. Liu, "The weight distribution of a family of p-ary cyclic codes," http://arxiv.org/pdf/1405.5278v1.pdf.

[22] X. Zeng, J. Shan, L. Hu, "A triple-error-correcting cyclic code from the Gold and Kasami-Welch APN power functions," *Finite Fields Appl.* **18**(1), 70-92(2012).

[23] X. Zeng, L. Hu, W. Jiang, Q. Yue, X. Cao, "The weight distribution of a class of p-ary cyclic codes," *Finite Fields Appl.* **16**(1), 56-73(2010).

[24] D. Zheng, X. Wang, X. Zeng, L. Hu, "The weight distribution of a family of p-ary cyclic codes," *Des. Codes Cryptogr.* doi: 10.1007/s10623-013-9908-2.

[25] Z. Zhou, C. Ding, "A class of three-weight cyclic codes," *Finite Fields Appl.* **25**, 79-93(2014).

[26] Z. Zhou, C. Ding, J. Luo, A. Zhang, "A family of five-weight cyclic codes and their weight enumerators," *IEEE Trans. Inform. Theory*, **59**(10), 6674-6682(2013).